

ANALYTICAL METHOD VALIDATION REPORT FOR ASSAY AND UNIFORMITY OF CONTENT OF RAMITHIAZIDE (RAMIPRIL AND HYDROCHLOROTHIAZIDE TABLETS) BY HPLC METHOD

Standard Test Wiethod Code: NA	Method Release Date: NA	
Version: 00	Supersedes Version: NA	
Reference: ICH guideline	Type of Validation: Validation of Non-Pharmacopoeial Method	
Customer Name: Generic Healthcare Pvt. Ltd.		

Prepared By	Designations	Signature	Date
Sachin Pujani	Manager Analytical (QbD research & Development Lab)	grijan	10-Aug-17
Reviewed By	Designation	Signature	Date
Satish Sawant	GM Analytical (QbD research & Development Lab)		10-Aug-17
Approved By	Designation	Signature	Date
Nargis penkan	QA Manager (QbD research & Development Lab)	Les ,	10-Aug-17
Approved By	Designation	Signature	Date
prabhakar Rawt	For (Generic Healthcare Pvt. Ltd.)	douter	10.08.2017



ANALYTICAL METHOD VALIDATION REPORT FOR ASSAY AND UNIFORMITY OF CONTENT OF RAMITHIAZIDE (RAMIPRIL AND HYDROCHLOROTHIAZIDE TABLETS) BY HPLC METHOD

Report Number QBD-REP-332 Page No. 2 of 49

INDEX

1.	OBJECTIVE	4
2.	SCOPE	4
3.	RESPONSIBILITIES	4
4.	SAFETY	4
5.	STANDARD DETAILS	4
6.	SAMPLE DETAILS	4
7.	REAGENTS AND CHEMICALS	5
8.	INSTRUMENT DETAILS	5
9.	METHOD OF ANALYSIS (FOR ASSAY AND CONTENT UNIFORMITY)	
10.	VALIDATION PARAMETERS	
10.1		
	SPECIFICITY	13
10.2		13 15
10.2 10.3	SPECIFICITY BY FORCED DEGRADATION:	15
10.3	SPECIFICITY BY FORCED DEGRADATION: LINEARITY AND RANGE	15 17
	2. SPECIFICITY BY FORCED DEGRADATION: 3. LINEARITY AND RANGE 4. PRECISION	15 17 21
10.3 10.4 10.5	2. SPECIFICITY BY FORCED DEGRADATION: 3. LINEARITY AND RANGE 4. PRECISION 5. ACCURACY (RECOVERY)	15 17 21 41
10.3 10.4	2. SPECIFICITY BY FORCED DEGRADATION: 3. LINEARITY AND RANGE 4. PRECISION 5. ACCURACY (RECOVERY) 6. ROBUSTNESS	15 21 41 43
10.3 10.4 10.5 10.6	2. SPECIFICITY BY FORCED DEGRADATION: 3. LINEARITY AND RANGE 4. PRECISION	15 21 41 43
10.3 10.4 10.5 10.6 10.7	2. SPECIFICITY BY FORCED DEGRADATION: 3. LINEARITY AND RANGE 4. PRECISION 5. ACCURACY (RECOVERY) 6. ROBUSTNESS 7. FILTER STUDY	15 21 41 43 45
10.3 10.4 10.5 10.6 10.7 10.8	2. SPECIFICITY BY FORCED DEGRADATION: 3. LINEARITY AND RANGE 4. PRECISION	15 21 41 43 45 49
10.3 10.4 10.5 10.6 10.7 10.8	2. SPECIFICITY BY FORCED DEGRADATION: 3. LINEARITY AND RANGE 4. PRECISION	15 21 41 43 45 49



ANALYTICAL METHOD VALIDATION REPORT FOR ASSAY AND UNIFORMITY OF CONTENT OF RAMITHIAZIDE (RAMIPRIL AND HYDROCHLOROTHIAZIDE TABLETS) BY HPLC METHOD

Report Number QBD-REP-332 Page No. 3 of 49

Abbreviations

	Abbleviations
mg	: Milligram
A.R. Grade	: Analytical Reagent Grade
μ	Micron
mL	: Milliliter
° C	: Degree Celsius
μΙ	: Microliter
G	: Gram
ppm	: Part Per Million
Std	: Standard
Wt.	: Weight
RSD	: Relative Standard Deviation
HPLC	: High Performance Liquid Chromatography
Conc.	: Concentration
NMT	: Not More Than
NLT	: Not Less Than
QA	: Quality Assurance



ANALYTICAL METHOD VALIDATION REPORT FOR ASSAY AND UNIFORMITY OF CONTENT OF RAMITHIAZIDE (RAMIPRIL AND HYDROCHLOROTHIAZIDE TABLETS) BY HPLC METHOD

Report Number	QBD-REP-332	Page No.	4 of 49

1. OBJECTIVE

The objective of the analytical method validation for Assay and Uniformity of content of Ramithiazide Tablets is to demonstrate, that the method will consistently yield results that will accurately reflect the quality characteristics of the product.

2. SCOPE

The validation activity has been carried out on Ramithiazide 5 mg/12.5 mg, Ramithiazide 10 mg/12.5 mg and Ramithiazide 10 mg/25 mg.

3. RESPONSIBILITIES

- 3.1 It is responsibility of the Manager Analytical, to prepare the Report.
- 3.2 It is the responsibility of the General Manager to review the Report.
- 3.3 It is the responsibility of the Manager QA to approve the Report.

4. SAFETY

Laboratory coats, safety gloves and goggles worn during this procedure and while handling chemicals. Normal laboratory codes should be followed and appropriate procedures adopted for disposal of waste.

5. STANDARD DETAILS

Standard Name	Batch No./Lot No.
Ramipril working standard	RL-14203
Hydrochlorothiazide working standard	20140403

^{*} Will be mentioned at the time of validation.

6. SAMPLE DETAILS

Sample Name	Batch No./Lot No.
Ramithiazide 5 mg/12.5 mg	GP170401
Ramithiazide 10 mg/12.5 mg	GP170402
Ramithiazide 10 mg/25 mg	GP170403
Placebo of Ramithiazide 5 mg/12.5 mg	-
Placebo of Ramithiazide 10 mg/12.5 mg	-
Placebo of Ramithiazide 10 mg/25 mg	-



ANALYTICAL METHOD VALIDATION REPORT FOR ASSAY AND UNIFORMITY OF CONTENT OF RAMITHIAZIDE (RAMIPRIL AND HYDROCHLOROTHIAZIDE TABLETS) BY HPLC METHOD

Report Number	QBD-REP-332	Page No.	5 of 49
report i amou	455 rai 552	2 008 2 1 00	

7. REAGENTS AND CHEMICALS

Reagent	Grade	Manufacturer
Sodium Perchlorate	AR	Finar
Acetonitrile	HPLC	Finar
Water	MilliQ	Inhouse

8. INSTRUMENT DETAILS

Instrument/ Equipment	Supplier/ Manufacturer / Make	Instrument/ Equipment In- House No.	Calibration Due Date
HPLC	Agilent	QbD-INST-HPLC-51	AUG-17
HPLC	Thermo	QbD-INST-HPLC-67	DEC-17
HPLC	Shimadzu	QbD-INST-HPLC-42	SEP-17
HPLC	Agilent	QbD-INST-HPLC-04	JAN-18
HPLC	Agilent	QbD-INST-HPLC-05	JAN-18
Dissolution Tester	Labindia	QbD-INST-DISS-09	OCT-17
Dissolution Tester	Labindia	QbD-INST-DISS-10	OCT-17
Analytical Balance	Mettler Toledo	QbD-INST-BAL-12	Daily
Analytical Balance	Mettler Toledo	QbD-INST-BAL-13	Daily
pH meter	Lab India	QbD-INST-pH-08	Daily

9. METHOD OF ANALYSIS (FOR ASSAY AND CONTENT UNIFORMITY)

HPLC Chromatographic Conditions:

Column	:	Sunniest C8, 150 mm x 4.6 mm,5 µm or equivalent
Flow rate	:	1.0 mL/min
Column temperature	:	25°C
LC mode	:	Isocratic
Detection wavelength	:	UV 210 nm
Injection Volume	:	10 μL
Run time	:	10 min

Diluent:

A mixture of 55 volumes of water, 45 volumes of Acetonitrile and 0.1 volume of triethylamine. Adjust pH 3.0 with dilute phosphoric acid.



ANALYTICAL METHOD VALIDATION REPORT FOR ASSAY AND UNIFORMITY OF CONTENT OF RAMITHIAZIDE (RAMIPRIL AND HYDROCHLOROTHIAZIDE TABLETS) BY HPLC METHOD

Report Number	QBD-REP-332	Page No.	6 of 49	
Keport Number	QDD-REI -332	Tage 110.	0 01 17	

Blank:

Use diluent as blank.

Preparation of buffer:

Dissolve 12.2 g of Sodium perchlorate in 1000 mL water and adjust the pH to 2.5 with diluted orthophosphoric acid. Filter through 0.45µ membrane filter and degas.

Preparation of mobile phase:

Mix 500 mL of buffer solution and 500 mL of acetonitrile.

Ramipril standard stock solution:

Weigh accurately and transfer 25.0 mg of Ramipril working standard into 25 mL of volumetric flask, add 10 mL of diluent, sonicate to dissolve and dilute to 25 mL with diluent and mix well. (Conc.: 1 mg/mL).

Hydrochlorothiazide standard stock solution:

Weigh accurately and transfer 25.0 mg of Hydrochlorothiazide working standard into 25 mL of volumetric flask, add 10 mL of diluent, sonicate to dissolve and dilute to 25 mL with diluent and mix well.(Conc.: 1 mg/mL).

Standard solution (For Ramithiazide Tablets 5 mg/12.5 mg and 10 mg/25 mg):

Dilute 2 mL of Ramipril standard stock solution and 5 mL of Hydrochlorothiazide standard stock solution to 100 mL with diluent and mix well. (Conc. 0.02 mg/mL of Ramipril and 0.05 mg/mL of Hydrochlorothiazide).

Standard solution (For Ramithiazide Tablets 10 mg/12.5 mg):

Dilute 2 mL of Ramipril standard stock solution and 2.5 mL of Hydrochlorothiazide standard stock solution to 100 mL with diluent and mix well. (Conc. 0.02 mg/mL of Ramipril and 0.025 mg/mL of Hydrochlorothiazide).



ANALYTICAL METHOD VALIDATION REPORT FOR ASSAY AND UNIFORMITY OF CONTENT OF RAMITHIAZIDE (RAMIPRIL AND HYDROCHLOROTHIAZIDE TABLETS) BY HPLC METHOD

Report Number QBD-REP-332 Page No. | 7 of 49

Sample solution (For Ramithiazide Tablets 5 mg/12.5 mg and 10 mg/25 mg):

Weight and powder 20 tablets. Weight accurately a quantity of powder containing about 25 mg of Ramipril, disperse in 5 mL of water, add 150 mL of diluent and sonicate for 15 min with intermittent shaking. Cool, dilute to 250 mL with diluent and mix well. Filter the solution through 0.45 μ nylon syringe filter. Dilute 5 mL of this solution to 25 mL with diluent and mix well. (Conc. 0.02 mg/mL of Ramipril and 0.05 mg/mL of Hydrochlorothiazide).

Sample solution (For Ramithiazide Tablets 10 mg/12.5 mg):

Weight and powder 20 tablets. Weight accurately a quantity of powder containing about 25 mg of Ramipril, disperse in 5 mL of water, add 150 mL of diluent and sonicate for 15 min with intermittent shaking. Cool, dilute to 250 mL with diluent and mix well. Filter the solution through 0.45 μ nylon syringe filter. Dilute 5 mL of this solution to 25 mL with diluent and mix well. (Conc. 0.2 mg/mL of Ramipril and 0.025 mg/mL of Hydrochlorothiazide).

Procedure:

Separately inject $10 \mu L$ of the blank solution, standard solution in five replicate and sample solution in duplicate into the chromatographic system, record the chromatograph and measure the peak response for the major peaks.

System suitability parameters:

The relative standard deviation for five replicate injections of standard solution should be not more than 2.0%. The theoretical plates of the principal peaks should be not less than 2000 and the tailing factor should be not more than 2.0.



ANALYTICAL METHOD VALIDATION REPORT FOR ASSAY AND UNIFORMITY OF CONTENT OF RAMITHIAZIDE (RAMIPRIL AND HYDROCHLOROTHIAZIDE TABLETS) BY HPLC METHOD

Report Number QBD-REP-332 Page No. 8 of 49

Calculations:

For Ramithiazide Tablets 5 mg/12.5 mg and 10 mg/25 mg

% Content of Ramipril

% Content =
$$AT WS 2 250 25 P$$

 $AS 25 100 WT 5 LC$

%Content of Hydrochlorothiazide

% Content =
$$AT1 WS1 5 250 25 P1$$

 $AS1 25 100 WT1 5 LC1$

For Ramithiazide Tablets 10 mg/12.5 mg

% Content of Ramipril

%Content of Hydrochlorothiazide

Where,

AT = Area of Ramipril peak obtained from sample preparation.

AS = Average area of Ramipril peak obtained from standard solution.

WS = Weight of Ramipril working standard in mg.

WT = Weight of sample in mg.

P = % Potency of Ramipril working standard on as is basis.

LC = Label claim of Ramipril per tablet in mg

AT1 = Area of Hydrochlorothiazide peak obtained from sample preparation.



ANALYTICAL METHOD VALIDATION REPORT FOR ASSAY AND UNIFORMITY OF CONTENT OF RAMITHIAZIDE (RAMIPRIL AND HYDROCHLOROTHIAZIDE TABLETS) BY HPLC METHOD

Report Number QBD-REP-332 Page No. 9 of 49

AS1 = Average area of Hydrochlorothiazide peak obtained from standard solution.

WS1 = Weight of Hydrochlorothiazide working standard in mg.

WT1 = Weight of sample in mg.

P1 = % Potency of Hydrochlorothiazide working standard on as is basis.

LC1 = Label claim of Hydrochlorothiazide per tablet in mg

UNIFORMITY OF CONTENT:

Note: HPLC Chromatographic conditions, Diluent, mobile phase, preparation of standard solutions are same as mention in assay method.

Sample solution (For Ramithiazide Tablets 5 mg/12.5 mg):

Add intact tablet into 250 mL volumetric flask. Disperse in 5 mL of water, add 150 mL of diluent and sonicate for 20 min with intermittent shaking. Cool and dilute to 250 mL with diluent and mix well. Filter the solution through 0.45 μ nylon syringe filter. (Conc. 0.2 mg/mL of Ramipril and 0.05 mg/mL of Hydrochlorothiazide).

Sample solution (For Ramithiazide Tablets 10 mg/12.5 mg):

Add intact tablet into 250 mL volumetric flask. Disperse in 5 mL of water, add 150 mL of diluent and sonicate for 20 min with intermittent shaking. Cool and dilute to 250 mL with diluent and mix well. Filter the solution through 0.45 μ nylon syringe filter. Dilute 5 mL of this solution to 10 mL with diluent and mix well. (Conc. 0.2 mg/mL of Ramipril and 0.025 mg/mL of Hydrochlorothiazide).

Sample solution (For Ramithiazide Tablets 10 mg/25 mg):

Add intact tablet into 250 mL volumetric flask. Disperse in 5 mL of water, add 150 mL of diluent and sonicate for 20 min with intermittent shaking. Cool and dilute to 250 mL with diluent and mix



ANALYTICAL METHOD VALIDATION REPORT FOR ASSAY AND UNIFORMITY OF CONTENT OF RAMITHIAZIDE (RAMIPRIL AND HYDROCHLOROTHIAZIDE TABLETS) BY HPLC METHOD

Report Number	QBD-REP-332	Page No.	10 of 49

well. Filter the solution through 0.45 μ nylon syringe filter. Dilute 10.0 mL of this solution to 20 mL with diluent and mix well. (Conc. 0.2 mg/mL of Ramipril and 0.05 mg/mL of Hydrochlorothiazide).

Procedure:

Separately inject $10~\mu L$ of the blank solution, standard solution in six replicate and ten sample solutions of uniformity of content into the chromatographic system, record the chromatograph and measure the peak response for the major peaks.

System suitability parameters:

The relative standard deviation for five replicate injections of standard solution should be not more than 2.0%. The theoretical plates of the principal peaks should be not less than 2000 and the tailing factor should be not more than 2.0.

Calculations:

For Ramithiazide Tablets 5 mg/12.5 mg

% Content of Ramipril =
$$AT WS 2 250 P$$

 $AS 25 100 1 LC$
% Content of HCTZ = $AT1 WS1 5 250 P1$
 $AS1 25 100 1 LC1$

For Ramithiazide Tablets 10 mg/12.5 mg

% Content of Ramipril =
$$AT$$
 WS 2 250 10 P

AS 25 100 1 5 LC

AT1 WS1 2.5 250 10 P1

% Content of HCTZ = $AT1$ WS1 2.5 250 10 P1

AS1 25 100 1 5 LC1



ANALYTICAL METHOD VALIDATION REPORT FOR ASSAY AND UNIFORMITY OF CONTENT OF RAMITHIAZIDE (RAMIPRIL AND HYDROCHLOROTHIAZIDE TABLETS) BY HPLC METHOD

Report NumberQBD-REP-332Page No.11 of 49

For Ramithiazide Tablets 10 mg/25 mg

% Content of Ramipril =
$$AT WS 2 250 10 P$$

 $AS 25 100 1 5 LC$

ATI WSI 5 250 10 PI

% Content of HCTZ = $ATI WSI 5 250 10 PI$
 $ASI 25 100 1 5 LCI$

Where,

AT = Area of Ramipril peak obtained from sample preparation.

AS = Average area of Ramipril peak obtained from standard solution.

WS = Weight of Ramipril working standard in mg.

WT = Weight of sample in mg.

P = % Potency of Ramipril working standard on as is basis.

LC = Label claim of Ramipril per tablet in mg

AT1 = Area of Hydrochlorothiazide peak obtained from sample preparation.

AS1 = Average area of Hydrochlorothiazide peak obtained from standard solution.

WS1 = Weight of Hydrochlorothiazide working standard in mg.

WT1 = Weight of sample in mg.

P1 = % Potency of Hydrochlorothiazide working standard on as is basis.

LC1 = Label claim of Hydrochlorothiazide per tablet in mg



ANALYTICAL METHOD VALIDATION REPORT FOR ASSAY AND UNIFORMITY OF CONTENT OF RAMITHIAZIDE (RAMIPRIL AND HYDROCHLOROTHIAZIDE TABLETS) BY HPLC METHOD

Report Number QBD-REP-332 Page No. 12 of 49

10. VALIDATION PARAMETERS

- 10.1. Specificity
- 10.2. Specificity by forced degradation
- 10.3. Linearity and Range
- 10.4. Precision
 - A. System precision
 - B. Method precision
 - C. Intermediate precision (Ruggedness)
- 10.5. Accuracy
- 10.6. Robustness
- 10.7. Filter study
- 10.8. Solution stability



ANALYTICAL METHOD VALIDATION REPORT FOR ASSAY AND UNIFORMITY OF CONTENT OF RAMITHIAZIDE (RAMIPRIL AND HYDROCHLOROTHIAZIDE TABLETS) BY HPLC METHOD

Report Number	QBD-REP-332	Page No.	13 of 49

10.1. SPECIFICITY

Specificity is the ability to asses unequivocally the analyte in the presence of components which may be expected to be present.

Acceptance Criteria:

- 1. The relative standard deviation for peak area of Ramipril and Hydrochlorothiazide of standard solution should be not more than 2.0%
- 2. The tailing factor for Ramipril and Hydrochlorothiazide peak in standard solution should be not more than 2.0
- 3. The theoretical plates for Ramipril and Hydrochlorothiazide peak in standard solution should be not less than 2000

Acceptance Criteria for Specificity:

There should not be any interference of peak due to blank and placebo at the retention time of the Ramipril and Hydrochlorothiazide peak.

Observations/Results:

Standard Details:

Injection Retention No. Time		Area of Hydrochlorothiazid Standard	e Reten		Area of Ramipril Standard				
1	2.10	27140.044	3.63	5	6928.843				
2	2.10	27182.012	3.63	5	6943.844				
3	2.10	27163.638	3.63	5	6935.208				
4	2.10	27160.989	3.63	5	6940.622 6923.226				
5	2.10	27129.962	3.63	3.65					
6	2.10	27176.337	3.63	5	6945.313				
Average	2.10	27158.830	3.63	5	6936.176				
SD	0.000	20.29	0.00	0	8.77				
%RSD 0.00		0.07	0.00	0.00 0.13					
	System Suitability Results								
% RSD of	Area of Hydro	% RSD of Area of Hydrochlorothiazide 0.07 NMT 2.0%							



ANALYTICAL METHOD VALIDATION REPORT FOR ASSAY AND UNIFORMITY OF CONTENT OF RAMITHIAZIDE (RAMIPRIL AND HYDROCHLOROTHIAZIDE TABLETS) BY HPLC METHOD

Report Number QBD-REP-332	Page No.	14 of 49
---------------------------	----------	----------

	% RSD of Area of Ramipril	0.13	NMT 2.0%
Theoretical plates of Ramipril		10410	NMT 2.0
The	oretical plates of Hydrochlorothiazide	10705	NMT 2.0
	Tailing factor of Ramipril	1.00	NLT 2000
Та	niling factor of Hydrochlorothiazide	1.04	NLT 2000

Interference Details:

Sample	Sample Weight (mg)	Retention Time (min)	Area of Hydrochlorothi azide	Retention Time (min)	Area of Ramipril
Blank Solution	NA	Not detected	Not detected	Not detected	Not detected
Standard Solution	Ramipril 25.9mg and 25.0mg of Hydrochlorot hiazide	2.10	27140.044	3.65	6928.843
Ramipril Identification Solution	25.9mg	3.65	6928.843	3.65	6910.967
Hydrochlorothiazide Identification Solution	25.0mg	2.10	26857.995	Not detected	Not detected
Placebo Solution	Not detected	Not detected	Not detected	Not detected	Not detected
Sample Solution	486.6	2.10	26811.141	3.65	6690.018

Conclusion:

No interference found due to blank and placebo at the retention time of the Hydrochlorothiazide and Ramipril.



ANALYTICAL METHOD VALIDATION REPORT FOR ASSAY AND UNIFORMITY OF CONTENT OF RAMITHIAZIDE (RAMIPRIL AND HYDROCHLOROTHIAZIDE TABLETS) BY HPLC METHOD

Report Number	QBD-REP-332	Page No.	15 of 49
reportrumer			

10.2. SPECIFICITY BY FORCED DEGRADATION:

Observations/Results:

Standard Details:

Injection No.	Retention Time	Area of Hydrochlorothia Standard	zide	Reten Tin		Area of Ramipril Standard
1	2.15	26533.941		4.1	5	7131.027
2	2.14	26555.379		4.1	5	7078.064
3	2.14	26620.305		4.1	5	7124.869
4	2.14	26611.268		4.1	5	7131.529
5	3.31	26610.794		5.3	2	7109.973
6	2.14	26650.629		4.15		7131.090
Average	2.34	26597.053		4.3	4	7117.759
SD	0.48	43.64		0.4	8	21.12
%RSD	20.42	0.16		10.9	99	0.30
		System Suitabi	lity R	esults		
% RSD of Are	a of Hydroch	lorothiazide	0	.16		NMT 2.0%
% RSD of Are	a of Ramipril		0	.30	NMT 2.0%	
Tailing factor	of Hydrochlor	rothiazide	1	.06		NMT 2.0
Tailing factor of Ramipril			1	.09		NMT 2.0
Theoretical plates of Hydrochlorothiazide				235		NLT 2000
Theoretical pla	ites of Ramip	ril	6.	328		NLT 2000



ANALYTICAL METHOD VALIDATION REPORT FOR ASSAY AND UNIFORMITY OF CONTENT OF RAMITHIAZIDE (RAMIPRIL AND HYDROCHLOROTHIAZIDE TABLETS) BY HPLC METHOD

Report Number	QBD-REP-332	Page No.	16 of 49

Sample Degradation Details (For Hydrochlorothiazide): Observations/Results:

Sample Type	Sample	Area of	% Assay	%	Peak
Untreated sample	494.71	26437.902	100.6	NA	999
Acid treated sample (5 mL of 1N HCl, 20 min at 70°C)	493.79	26539.404	101.2	-0.6	999
Base treated sample (5 mL of 1N NaOH, 20 min at 70°C)	493.79	26555.589	101.2	-0.6	999
Hydrogen peroxide treated sample (5 mL of 30%H2O2,	494.31	26541.634	101.1	-0.5	999
Acid treated sample (2 mL of 1N HCl, 29 min at 70°C)	493.96	26502.271	101.0	-0.4	999
Base treated sample (2 mL of 1N NaOH, 20 min at 70°C)	494.26	26475.967	100.8	-0.2	999
Sample solution exposed to Heat	493.70	26918.472	102.6	-2.0	999

Sample Degradation Details (For Ramipril):

Sample Type	Sample Weight (mg)	Area of Sample	% Assay	% Degradation	Peak purity Match
Untreated sample	494.71	6990.951	99.3	NA	999
Acid treated sample (5 mL of 1N HCl, 20 min at 70°C)	493.79	6975.559	99.3	0.0	999
Base treated sample (5 mL of 1N NaOH, 20 min at	493.79	6896.102	98.2	1.1	999
Hydrogen peroxide treated sample (5 mL of	494.31	6921.174	98.4	0.9	999
Acid treated sample (2 mL of 1N HCl, 20 min at 70°C)	493.96	6963.870	99.1	0.2	999
Base treated sample (2 mL of 1N NaOH, 20 min at	494.26	6918.869	98.4	0.9	999
Sample solution exposed to Heat	493.70	7082.866	100.8	-1.5	999

Conclusion:

From the above data it is observed that product is not degraded in above conditions.



ANALYTICAL METHOD VALIDATION REPORT FOR ASSAY AND UNIFORMITY OF CONTENT OF RAMITHIAZIDE (RAMIPRIL AND HYDROCHLOROTHIAZIDE TABLETS) BY HPLC METHOD

Report Number	QBD-REP-332	Page No.	17 of 49

10.3. LINEARITY AND RANGE

The linearity of an analytical procedure is its ability (within a given range) to obtain test results which are directly proportional to the concentration (amount) of analyte in the sample.

The range of an analytical procedure is the interval between the upper and lower concentration (amounts) of analyte in the sample (including these concentrations) for which it has been demonstrated that the analytical procedure has a suitable level of precision, accuracy and linearity.

Acceptance Criteria:

- 1. The relative standard deviation for peak area of Ramipril and Hydrochlorothiazide of standard solution should be not more than 2.0%
- 2. The tailing factor for Ramipril and Hydrochlorothiazide peak in standard solution should be not more than 2.0
- 3. The theoretical plates for Ramipril and Hydrochlorothiazide peak in standard solution should be not less than 2000
- 4. The correlation coefficient obtained from the graph should be NLT 0.99

Observations/Results:

Standard Details:

Injection No.	Retention Time	Area of Hydrochlorothiaz Standard	zide	Retent Tim		Area of Ramipril Standard
1	2.113	27192.801		3.75	3	7007.386
2	2.113	27239.583		3.75	3	7018.219
3	2.117	27322.059		3.75	3	7031.934
4	2.117	27238.009		3.75	7	7008.798
5	2.117	27175.641		3.75	7	6989.347
6	2.113	27184.405		3.75	3	6992.041
Average	2.115	27225.416		3.75	4	7007.954
SD	0.00	54.63		0.00)	16.01
%RSD	0.10	0.20		0.06)	0.23
System Suitability Results						
% RSD of Are	% RSD of Area of Hydrochlorothiazide			.20		NMT 2.0%
% RSD of Are	a of Ramipril		0	.23		NMT 2.0%



ANALYTICAL METHOD VALIDATION REPORT FOR ASSAY AND UNIFORMITY OF CONTENT OF RAMITHIAZIDE (RAMIPRIL AND HYDROCHLOROTHIAZIDE TABLETS) BY HPLC METHOD

Report Number	QBD-REP-332	Page No.	18 of 49
---------------	-------------	----------	----------

Tailing factor of Hydrochlorothiazide	1.09	NMT 2.0
Tailing factor of Ramipril	1.00	NMT 2.0
Theoretical plates of Hydrochlorothiazide	10362	NLT 2000
Theoretical plates of Ramipril	9677	NLT 2000


Linearity Determination for Hydrochlorothiazide:

Level	Linearity level (%)	ml of Standard Stock Solution added	Diluted to volume (mL)	Actual Concentration (ppm)	Area of sample	Average area	
					6759.543		
					6742.148		
1	25	1.2	100	12.1	6761.846	6761.625	
1	23	1.2	100		6740.715	0701.023	
					6798.745		
					6766.757		
					13907.523		
2	50	2.5	100	25.3	13902.865	13905.135	
					13905.017		
					22197.151		
3	75	4.0	100	40.4	22158.106	22179.585	
					22183.498		
					27155.943		
4	100	5.0	100	50.5	27084.756	27122.050	
					27125.449		
					32725.509		
5	125	6.0	100	60.6	32723.357	32723.721	
				,	32722.296		
					40376.322		
			40480.045				
6	150	7.5	100	75.8	40458.462	10200 540	
6 150 7.5	100	13.0	40360.603	40390.549			
					40379.248		
					40288.613		



ANALYTICAL METHOD VALIDATION REPORT FOR ASSAY AND UNIFORMITY OF CONTENT OF RAMITHIAZIDE (RAMIPRIL AND HYDROCHLOROTHIAZIDE TABLETS) BY HPLC METHOD

Report Number QBD-REP-332 Page No. 19 of 49



Slope	528.69
y-Intercept	530.01
Correleation Coefficient	0.9999

Linearity Determination for Ramipril:


Level	Linearity level (%)	ml of Standard Stock Solution added	Diluted to volume (mL)	Actual Concentration (ppm)	Area of sample	Average area		
					1793.527			
					1792.789			
1	25	0.5	0.5	1 100 50	5.0	100 5.0	1785.713	1789.150
1	25						1780.852	
					1795.589			
					1786.431			
					3531.387			
2	50	1.0	100	10.1	3535.378	3531.410		
					3527.465			
					5672.921			
3	75	1.6	100	16.1	5674.689	5674.802		
					5676.795			
					6998.638			
4	100	2.0	100	100	20.1	6976.426	6984.227	
					6977.616			



ANALYTICAL METHOD VALIDATION REPORT FOR ASSAY AND UNIFORMITY OF CONTENT OF RAMITHIAZIDE (RAMIPRIL AND HYDROCHLOROTHIAZIDE TABLETS) BY HPLC METHOD

Report Number (QBD-REP-332	Page No.	20 of 49
-----------------	-------------	----------	----------

5	125	2.4	100	24.2	8484.320 8476.854 8489.456	8483.543
6	150	3.0	100	30.2	10516.722 10531.706 10542.009 10499.757 10505.742 10510.904	10517.807



Slope	347.37
y-Intercept	43.12
Correleation Coefficient	0.9999

Conclusion:

- 1. The relative standard deviation for peak area of Ramipril and Hydrochlorothiazide of standard solution is within acceptance criteria.
- 2. The tailing factor for Ramipril and Hydrochlorothiazide peak in standard solution is within acceptance criteria.
- 3. The theoretical plates for Ramipril and Hydrochlorothiazide peak in standard solution is within acceptance criteria.
- 4. The correlation coefficient obtained from the graph for Hydrochlorothiazide is 0.9999 and for Ramipril is 0.9999.



ANALYTICAL METHOD VALIDATION REPORT FOR ASSAY AND UNIFORMITY OF CONTENT OF RAMITHIAZIDE (RAMIPRIL AND HYDROCHLOROTHIAZIDE TABLETS) BY HPLC METHOD

Report Number	QBD-REP-332	Page No.	21 of 49
Report Number	QDD-REI -332	Tage 110.	21 01 17

10.4. PRECISION

Precision of the method shall be established by carrying out as System precision, Method precision and Intermediate precision.

A) System Precision

Acceptance Criteria:

- 1. The relative standard deviation for peak area of Ramipril and Hydrochlorothiazide of standard solution should be not more than 2.0%
- 2. The tailing factor for Ramipril and Hydrochlorothiazide peak in standard solution should be not more than 2.0
- 3. The theoretical plates for Ramipril and Hydrochlorothiazide peak in standard solution should be not less than 2000

Observations/Results:

Standard Details (Ramipril 5 mg & Hydrochlorothiazide 12.5 mg & Ramipril 5 mg & Hydrochlorothiazide 12.5 mg)

Injection No.	Retention Time	Area of Hydrochlorothia Standard	zide	Retenti		Area of Ramipril Standard
1	2.10	27140.044		3.65		6928.843
2	2.10	27182.012		3.65		6943.844
3	2.10	27163.638		3.65		6935.208
4	2.10	27160.989		3.65		6940.622
5	2.10	27129.962		3.65		6923.226
6	2.10	27176.338		3.65		6945.313
Average	2.10	27158.830		3.65		6936.176
SD	0.000	20.29		0.000)	8.77
%RSD	0.00	0.07		0.00		0.13
		System Suitabi	lity R	esults		
% RSD of Rar	nipril	•	0	.13		NMT 2.0%
% RSD of Hyd	drochlorothiaz	zide	0	.07	7 NMT 2.0%	
Therotical plat	es of Ramipri	1	10	410 NMT 2.0		NMT 2.0
Therotical plates of Hydrochlorothiazide			10	705	705 NMT 2.0	
Tailing factor of Ramipril				1.00 NLT 200		NLT 2000
Tailing factor		lorothiazide	1	.04		NLT 2000



ANALYTICAL METHOD VALIDATION REPORT FOR ASSAY AND UNIFORMITY OF CONTENT OF RAMITHIAZIDE (RAMIPRIL AND HYDROCHLOROTHIAZIDE TABLETS) BY HPLC METHOD

Report Number	QBD-REP-332	Page No.	22 of 49

Standard Details (Ramipril 10 mg & Hydrochlorothiazide 12.5 mg)

Injection No.	Retention Time	Area of Hydrochlorothia Standard	zide	Reten Tin		Area of Ramipril Standard
1	2.10	14123.816		3.6	5	7005.022
2	2.10	14086.982		3.6	5	6985.895
3	2.10	14137.983		3.6	5	7006.536
4	2.10	14224.821		3.6	5	7039.733
5	2.10	14154.966		3.6	5	7024.769
6	2.10	14132.359		3.6	5	6996.007
Average	2.10	14143.488		3.6	5	7009.660
SD	0.000	45.78		0.00	00	19.57
%RSD	0.00	0.32		0.0	0	0.28
		System Suitabi	lity R	Results		
% RSD of Are	a of Hydroch	lorothiazide	0	.32		NMT 2.0%
% RSD of Are	a of Ramipril		0	.28	NMT 2.0%	
Tailing factor of Hydrochlorothiazide		10	768		NMT 2.0	
Tailing factor of Ramipril		10	338	NMT 2.0		
Theoretical pla	tes of Hydro	chlorothiazide	1	.06		NLT 2000
Theoretical pla	tes of Ramip	ril	0	.98		NLT 2000

Conclusion:

- 1. The relative standard deviation for peak area of Ramipril and Hydrochlorothiazide of standard solution is within acceptance criteria.
- 2. The tailing factor for Ramipril and Hydrochlorothiazide peak in standard solution is within acceptance criteria.
- 3. The theoretical plates for Ramipril and Hydrochlorothiazide peak in standard solution is within acceptance criteria.

B) Method Precision

The precision of an analytical procedure expresses the closeness of agreement (degree of scatter) between a series of measurements obtained from multiple sampling of the same homogeneous sample under the prescribed conditions.



ANALYTICAL METHOD VALIDATION REPORT FOR ASSAY AND UNIFORMITY OF CONTENT OF RAMITHIAZIDE (RAMIPRIL AND HYDROCHLOROTHIAZIDE TABLETS) BY HPLC METHOD

	ODD DED 222	D M	22 - 6.40
Report Number	QBD-REP-332	Page No.	23 01 49

I) Method Precision For Assay:

Acceptance Criteria:

- 1. The relative standard deviation for peak area of Ramipril and Hydrochlorothiazide of standard solution should be not more than 2.0%
- 2. The tailing factor for Ramipril and Hydrochlorothiazide peak in standard solution should be not more than 2.0
- 3. The theoretical plates for Ramipril and Hydrochlorothiazide peak in standard solution should be not less than 2000
- 4. % Assay value of Ramipril and Hydrochlorothiazide for sample solution should be within specified limit.
- 5. % RSD of Assay value of Ramipril and Hydrochlorothiazide for six sample solution should be not more than 2.0%

i) Precision Sample Details (Ramipril 5 mg & Hydrochlorothiazide 12.5 mg):

Sample	Sample			% Assay		
No.	vveigni	Hydrochlorothiazide	Ramipril	Hydrochlorothiazide	Ramipril	
1	975.1	26742.050	6512.104	100.8	99.5	
2	975.0	26540.983	6430.609	100.0	98.2	
3	974.5	26760.319	6552.363	100.9	100.1	
4	974.2	26787.668	6502.765	101.1	99.4	
5	975.1	26969.459	6559.458	101.7	100.2	
6	974.3	26818.478	6507.422	101.2	99.5	
			Mean	100.9	99.5	
			SD	0.53	0.71	
			%RSD	0.52	0.71	



ANALYTICAL METHOD VALIDATION REPORT FOR ASSAY AND UNIFORMITY OF CONTENT OF RAMITHIAZIDE (RAMIPRIL AND HYDROCHLOROTHIAZIDE TABLETS) BY HPLC METHOD

Report Number QBD-REP-332	Page No.	24 of 49
---------------------------	----------	----------

ii) Precision Sample Details (Ramipril 10 mg & Hydrochlorothiazide 12.5 mg):

Sample	Sample			% Assay		
No.	Weight (mg)	Hydrochlorothiazide	Ramipril	Hydrochlorothiazide	Ramipril	
1	498.5	13979.161	6902.898	98.2	101.3	
2	525.1	14777.095	7296.229	98.6	101.6	
3	508.5	14296.125	7057.916	98.5	101.5	
4	502.1	14110.117	6959.222	98.4	101.4	
5	501.3	14161.119	6957.830	98.9	101.5	
6	496.7	13971.434	6884.103	98.5	101.4	
			Average	98.5	101.5	
			SD	0.24	0.13	
			%RSD	0.24	0.13	

iii) Precision Sample Details (Ramipril 10 mg & Hydrochlorothiazide 25 mg):

Sample	Sample	1 11 000 01 200111		% Assay		
No.	Weight (mg)	Hydrochlorothiazide	Ramipril	Hydrochlorothiazide	Ramipril	
1	486.6	26811.141	6690.018	99.4	100.5	
2	483.8	26836.422	6700.093	100.1	101.2	
3	494.5	27483.766	6868.453	100.3	101.5	
4	491.9	26804.727	6712.113	98.3	99.7	
5	497.8	27713.304	6920.976	100.4	101.6	
6	489.2	26963.565	6753.191	99.4	100.9	
		'	Average	99.6	100.9	
			SD	0.79	0.71	
			%RSD	0.79	0.71	



ANALYTICAL METHOD VALIDATION REPORT FOR ASSAY AND UNIFORMITY OF CONTENT OF RAMITHIAZIDE (RAMIPRIL AND HYDROCHLOROTHIAZIDE TABLETS) BY HPLC METHOD

Report Number	QBD-REP-332	Page No.	25 of 49
---------------	-------------	----------	----------

Conclusion:

- 1. The relative standard deviation for peak area of Ramipril and Hydrochlorothiazide of standard solution should be not more than 2.0%
- 2. The tailing factor for Ramipril and Hydrochlorothiazide peak in standard solution is within acceptance criteria.
- 3. The theoretical plates for Ramipril and Hydrochlorothiazide peak in standard solution is within acceptance criteria.
- 4. % Assay value of Ramipril and Hydrochlorothiazide for sample solution is within acceptance criteria.
- 5. % RSD of Assay value of Ramipril and Hydrochlorothiazide for six sample solution is within acceptance criteria.

II) Method Precision for Content Uniformity:

Acceptance Criteria:

- 1. The relative standard deviation for peak area of Ramipril and Hydrochlorothiazide of standard solution should be not more than 2.0%
- 2. The tailing factor for Ramipril and Hydrochlorothiazide peak in standard solution should be not more than 2.0
- 3. The theoretical plates for Ramipril and Hydrochlorothiazide peak in standard solution should be not less than 2000
- 4. The acceptance value of Ramipril and Hydrochlorothiazide for ten sample solution should be not more than 15.0%



ANALYTICAL METHOD VALIDATION REPORT FOR ASSAY AND UNIFORMITY OF CONTENT OF RAMITHIAZIDE (RAMIPRIL AND HYDROCHLOROTHIAZIDE TABLETS) BY HPLC METHOD

Report Number	QBD-REP-332	Page No.	26 of 49
---------------	-------------	----------	----------

Standard Details (Ramipril 5 mg & Hydrochlorothiazide 12.5 mg & Ramipril 10 mg & Hydrochlorothiazide 25 mg)

Injection No.	Retention Time	Area of Hydrochlorothiazide Standard	Retentio Time	n Area of Ramipril Standard
1	2.11	26693.654	3.75	6998.101
2	2.11	26623.399	3.75	6978.537
3	2.11	26584.575	3.75	6964.311
4	2.11	26571.035	3.74	6962.451
5	2.11	26622.841	3.74	6983.342
6	2.11	26563.407	3.74	6955.304
Average	2.11	26609.819	3.74	6973.674
SD	0.00	48.33	0.01	15.91
%RSD	0.00	0.18	0.14	0.23
		System Suitability	Results	
% RSD of A	rea of Hydro	chlorothiazide	0.18	NMT 2.0%
% RSD of Area of Ramipril		ril	0.23	NMT 2.0%
Tailing factor of Hydrochlorothiazide		1.08	NMT 2.0	
Tailing factor of Ramipril		1.03	NMT 2.0	
Theoretical	plates of Hydi	rochlorothiazide	10583	NLT 2000
Theoretical	plates of Ram	ipril	11081	NLT 2000

Standard Details (Ramipril 10 mg & Hydrochlorothiazide 12.5 mg)

		Area of			
Injection No.	Retention Time	Hydrochlorothiazid Standard	le	Retenti Time	Area of Ramipril Standard
1	2.11	13589.204		3.73	7021.068
2	2.11	13576.131		3.73	7015.366
3	2.11	13614.588		3.73	7032.524
4	2.11	13634.116		3.73	7052.893
5	2.11	13630.656		3.73	7047.321
6	2.11	13610.292		3.73	7014.270
Average	2.11	13609.164		3.73	7030.573
SD	0.00	22.82		0.00	16.55
%RSD	0.00	0.17		0.00	0.24
		System Suitabilit	y R	esults	
% RSD of A	Area of Hydro	chlorothiazide		0.24	NMT 2.0%
% RSD of A	Area of Ramip	oril		0.17	NMT 2.0%
Tailing fact	or of Hydroch	lorothiazide		1.14	NMT 2.0
Tailing factor of Ramipril			1.04	NMT 2.0	
Theoretical	Theoretical plates of Hydrochlorothiazide		1	0841	NLT 2000
Theoretical	plates of Ran	nipril	1	1040	NLT 2000



ANALYTICAL METHOD VALIDATION REPORT FOR ASSAY AND UNIFORMITY OF CONTENT OF RAMITHIAZIDE (RAMIPRIL AND HYDROCHLOROTHIAZIDE TABLETS) BY HPLC METHOD

Report Number QBD-REP-332 Page No. 27 of 49

i) Precision sample details for Hydrochlorothiazide(Ramipril 5mg & Hydrochlorothiazide 12.5mg):

Sample	Tablets Weight (mg)	Peak Area of Sample Hydrochlorothiazide	% Content			
1	920.47	27629.543	103.9			
2	926.28	27305.449	102.7			
3	905.59	27466.866	103.3			
4	920.14	27673.941	104.1			
5	930.56	27967.369	105.2			
6	904.67	27599.831	103.8			
7	912.88	27678.173	104.1			
8	927.36	27388.930	103.0			
9	892.39	27595.452	103.8			
10	895.88	27537.136	103.6			
		Min	102.7			
		Max	105.2			
		Mean	103.8			
		SD	0.68			
		% RSD	0.66			
		AV Value	3.9			

ii) Precision sample details for Ramipril (Ramipril 5 mg & Hydrochlorothiazide 12.5 mg):

1 (the print ing et 11) di temor temazide 12.5 ing).					
Sample	Tablets Weight (mg)	Peak Area of Sample Ramipril	% Content		
1	920.47	6757.220	100.0		
2	926.28	6609.463	97.8		
3	905.59	6679.661	98.8		
4	920.14	6508.468	96.3		
5	930.56	6675.710	98.8		
6	904.67	6809.909	100.8		
7	912.88	6762.906	100.1		
8	927.36	6639.258	98.2		
9	892.39	6503.552	96.2		
10	895.88	6475.370	95.8		
		Min	95.8		
		Max	100.8		
		Mean	98.3		
		SD	1.74		
		% RSD	1.77		
		AV Value	1.8		



ANALYTICAL METHOD VALIDATION REPORT FOR ASSAY AND UNIFORMITY OF CONTENT OF RAMITHIAZIDE (RAMIPRIL AND HYDROCHLOROTHIAZIDE TABLETS) BY HPLC METHOD

Report NumberQBD-REP-332Page No.28 of 49

iii) Precision Sample Details for Hydrochlorothiazide (Ramipril 10 mg & Hydrochlorothiazide 12.5 mg):

Sample	Tablets Weight (mg)	Peak Area of Sample	% Content
1	920.47	13862.943	102.0
2	926.28	14268.605	105.0
3	905.59	14036.682	103.2
4	920.14	14088.776	103.6
5	930.56	14075.274	103.5
6	904.67	14061.901	103.4
7	912.88	14062.203	103.4
8	927.36	14006.217	103.0
9	892.39	14186.247	104.3
10	895.88	13900.783	102.2
		Min	102.0
		Max	105.0
		Mean	103.4
		SD	0.88
		% RSD	0.85
		AV Value	3.7

iv)Precision Sample Details for Ramipril (Ramipril 10 mg & Hydrochlorothiazide 12.5 mg):

Sample	Tablets Weight (mg)	Peak Area of Sample	% Content
1	920.47	6934.208	101.8
2	926.28	6830.801	100.3
3	905.59	6903.521	101.3
4	920.14	6837.118	100.4
5	930.56	6885.831	101.1
6	904.67	6841.661	100.4
7	912.88	6953.359	102.1
8	927.36	6896.770	101.2
9	892.39	6814.032	100.0
10	895.88	6888.926	101.1
		Min	100.0
		Max	102.1
		Mean	101.0
		SD	0.68
		% RSD	0.67
		AV Value	1.6



ANALYTICAL METHOD VALIDATION REPORT FOR ASSAY AND UNIFORMITY OF CONTENT OF RAMITHIAZIDE (RAMIPRIL AND HYDROCHLOROTHIAZIDE TABLETS) BY HPLC METHOD

Report Number	QBD-REP-332	Page No.	29 of 49

v) Precision sample details for Hydrochlorothiazide (Ramipril 10 mg & Hydrochlorothiazide 25 mg):

Sample	Tablets Weight (mg)	Peak Area of Sample	% Content
1	920.47	26198.918	98.6
2	926.28	27036.530	101.7
3	905.59	27303.232	102.7
4	920.14	27055.505	101.8
5	930.56	27397.083	103.1
6	904.67	27495.587	103.4
7	912.88	27296.913	102.7
8	927.36	27487.205	103.4
9	892.39	27138.350	102.1
10	895.88	27202.516	102.3
		Min	98.6
		Max	103.4
		Mean	102.2
		SD	1.41
		% RSD	1.38
		AV Value	1.6

vi)Precision sample details for Ramipril (Ramipril 10 mg & Hydrochlorothiazide 25.5 mg):

Sample	Tablets Weight (mg)	Peak Area of Sample	% Content
1	920.47	6248.850	92.5
2	926.28	6572.144	97.3
3	905.59	6775.708	100.3
4	920.14	6445.797	95.4
5	930.56	6596.430	97.6
6	904.67	6675.208	98.8
7	912.88	6676.491	98.8
8	927.36	6699.905	99.1
9	892.39	6531.341	96.6
10	895.88	6570.784	97.2
		Min	92.47
		Max	100.26
		Mean	97.4
		SD	2.21
		% RSD	2.27
		AV Value	7.5



ANALYTICAL METHOD VALIDATION REPORT FOR ASSAY AND UNIFORMITY OF CONTENT OF RAMITHIAZIDE (RAMIPRIL AND HYDROCHLOROTHIAZIDE TABLETS) BY HPLC METHOD

Report Number QBD-REP-332 Page No. 30 of 49

Conclusion:

- 1. The relative standard deviation for peak area of Ramipril and Hydrochlorothiazide of standard solution should be not more than 2.0%
- 2. The tailing factor for Ramipril and Hydrochlorothiazide peak in standard solution is within acceptance criteria.
- 3. The theoretical plates for Ramipril and Hydrochlorothiazide peak in standard solution is within acceptance criteria.
- 4. The acceptance value of Ramipril and Hydrochlorothiazide for ten sample solution is within acceptance criteria.

C) Intermediate Precision

Intermediate precision of the method will be established by carrying out the analysis on different day with another analyst and another instrument.

I) Method precision For Assay:

Acceptance criteria:

- 1. The relative standard deviation for peak area of Ramipril and Hydrochlorothiazide of standard solution should be not more than 2.0%
- 2. The tailing factor for Ramipril and Hydrochlorothiazide peak in standard solution should be not more than 2.0
- 3. The theoretical plates for Ramipril and Hydrochlorothiazide peak in standard solution should be not less than 2000
- 4. % Assay value of Ramipril and Hydrochlorothiazide for sample solution should be within specified limit.
- 5. % RSD of Assay value of Ramipril and Hydrochlorothiazide for six sample solution should be not more than 2.0%
- The cumulative RSD for % Assay of Ramipril and Hydrochlorothiazide for twelve samples in method precision and intermediate precision should be not more than 2.0%



ANALYTICAL METHOD VALIDATION REPORT FOR ASSAY AND UNIFORMITY OF CONTENT OF RAMITHIAZIDE (RAMIPRIL AND HYDROCHLOROTHIAZIDE TABLETS) BY HPLC METHOD

	Report Number	QBD-REP-332	Page No.	31 of 49
-			Tage 110.	31 01 17

Observations/Results:

Standard details (Ramipril 5 mg & Hydrochlorothiazide 12.5 mg & Ramipril 10 mg & Hydrochlorothiazide 25 mg)

Injection Details						
Injection	Retention	Area of		Rete	ntion	Area of Ramipril
No.	Time	Hydrochlorothia	zide	Tir	ne	Standard
1	2.225	29779.744		3.8	82	6982.083
2	2.225	29910.378		3.8	83	6990.418
3	2.227	29895.007		3.8	85	6982.654
4	2.227	29870.134		3.8	85	6978.374
5 2.225 29937.11		29937.184		3.8	83	6982.466
6	2.225	29970.544		3.8	83	6990.327
Average	2.226	29893.832		3.8	84	6984.387
SD	0.001	65.75		0.0	01	4.89
%RSD	0.05	0.22		0.0	3	0.07
		System Suitabi	lity R	esults		
% RSD of Area		orothiazide	_	22		NMT 2.0%
% RSD of Area	a of Ramipril		0.	07	11111 21070	
Tailing factor of Hydrochlorothiazide			940			
Tailing factor of Ramipril			84	NMT 2.0		
Theoretics plan	tes of Hydroc	hlorothiazide		15		NLT 2000
Theoretical plan				06		NLT 2000

Standard details (Ramipril 10 mg & Hydrochlorothiazide 12.5 mg)

Injection	Retention	Area of		Reter	tion	Area of Ramipril
1	2.223	15522.964		3.8	82	7108.906
2	2.225	15529.755		3.8		7106.701
3	2.225	15525.629		3.8	83	7101.858
4	2.223	15531.716		3.8		7102.031
5	2.223	15526.224		3.8	32	7098.623
6	2.223	15534.615		3.88	32	7101.488
Average	2.224	15528.484		3.88		
SD	0.001	4.32		0.00	00	3.79
%RSD	0.05	0.03		0.0	1	0.05
		System Suitabi	lity R	esults		
% RSD of Area		orothiazide	0.	.03	NMT 2.0%	
% RSD of Area			0.	.05		
Tailing factor of Hydrochlorothiazide		58	365			
Tailing factor of Ramipril		83	71	NMT 2.0		
Theoretical pla			1.	13		NLT 2000
Theoretical pla	tes of Ramipr	il	1.	06		NLT 2000



ANALYTICAL METHOD VALIDATION REPORT FOR ASSAY AND UNIFORMITY OF CONTENT OF RAMITHIAZIDE (RAMIPRIL AND HYDROCHLOROTHIAZIDE TABLETS) BY HPLC METHOD

Report Number	QBD-REP-332	Page No.	32 of 49

i) Intermediate precision sample details (Ramipril 5 mg & Hydrochlorothiazide 12.5 mg):

Sample Sample		Area of San	Area of Sample		% Assay		
No.	Weight (mg)	Hydrochlorothiazide	Ramipril	Hydrochlorothiazide	Ramipril		
1	995.72	29565.134	7130.849	99.8	100.4		
2	994.71	29626.454	7135.219	100.1	100.5		
3	995.09	29548.641	7138.142	99.8	100.5		
4	995.33	29467.116	7116.769	99.5	100.2		
5	995.73	29706.000	7160.037	100.3	100.8		
6	995.17	29503.873	7124.497	99.6	100.3		
			Mean	99.8	100.5		
			SD	0.29	0.20		
			%RSD	0.29	0.20		

ii) Intermediate precision sample details (Ramipril 10 mg & Hydrochlorothiazide 12.5 mg):

Sample No. Sample Weight (mg)		Area of Sample		% Assay		
		Hydrochlorothiazide	Ramipril	Hydrochlorothiazide	Ramipril	
1	494.49	15452.158	7121.486	100.3	98.5	
2	494.59	15470.739	7129.698	100.4	98.6	
3	494.95	15490.150	7139.473	100.5	98.7	
4	495.81	15432.785	7115.849	99.9	98.2	
5	495.66	15588.560	7154.534	101.0	98.7	
6	494.98	15574.824	7135.676	101.0	98.6	
			Average	100.5	98.5	
			SD	0.41	0.20	
			%RSD	0.41	0.20	



ANALYTICAL METHOD VALIDATION REPORT FOR ASSAY AND UNIFORMITY OF CONTENT OF RAMITHIAZIDE (RAMIPRIL AND HYDROCHLOROTHIAZIDE TABLETS) BY HPLC METHOD

Report Number QBD-REP-332 Page No. 33 of 49

iii) Intermediate precision sample details (Ramipril 10 mg & Hydrochlorothiazide 25 mg):

Sample No.	Sample Weight (mg)	Area of Sam	ıple	% Assay			
		Hydrochlorothiazide	Ramipril	Hydrochlorothiazide	Ramipril		
1	493.22	29503.345	7122.542	98.7	99.3		
2	493.35	29483.484	7119.263	98.6	99.3		
3	493.63	29462.967	7116.801	98.4	99.2		
4	493.27	29499.358	7130.428	98.6	99.4		
5	493.44	29512.245	7118.175	98.6	99.2		
6	493.83	29534.488	7120.850	98.6	99.2		
			Average	98.6	99.3		
			SD	0.08	0.10		
			%RSD	0.08	0.10		

Comparison of Method Precision and Intermediate precision:

i) For Ramipril 5 mg & Hydrochlorothiazide 12.5 mg

Sample	Parameter	% Assay			
No.		Hydrochlorothiazide	Ramipril		
1		100.8	99.5		
2		100.0	98.2		
3	Method precision	100.9	100.1		
4		101.1	99.4		
5		101.7	100.2		
6		101.2	99.5		
1		99.8	100.4		
2		100.1	100.5		
3	Intermediate precision	99.8	100.5		
4	intermediate precision	99.5	100.2		
5		100.3	100.8		
6		99.6	100.3		
	Mean	100.4	100.0		
	SD	0.72	0.71		
	%RSD	0.71	0.71		



ANALYTICAL METHOD VALIDATION REPORT FOR ASSAY AND UNIFORMITY OF CONTENT OF RAMITHIAZIDE (RAMIPRIL AND HYDROCHLOROTHIAZIDE TABLETS) BY HPLC METHOD

Report Number	QBD-REP-332	Page No.	34 of 49
-		I age 110.	510177

ii) For Ramipril 10 mg & Hydrochlorothiazide 12.5 mg

Sample	Parameter	% Assay			
No.		Hydrochlorothiazide	Ramipril		
98.2		98.2	101.3		
98.6		98.6	101.6		
98.5	Method precision	98.5	101.5		
98.4		98.4	101.4		
98.9		98.9	101.5		
98.5		98.5	101.4		
100.3	98.5	100.3	98.5		
100.4	98.6	100.4	98.6		
100.5	98.7	100.5	98.7		
99.9	98.2	99.9	98.2		
101	98.7	101	98.7		
101	98.6	101	98.6		
99.5	100.0	99.5	100.0		
1.09	1.52	1.09	1.52		
1.10	1.52	1.10	1.52		

iii) For Ramipril 10 mg & Hydrochlorothiazide 25 mg

Sample	Parameter	% Assay			
No.		Hydrochlorothiazide	Ramipril		
1		99.4	100.5		
2	Method precision	100.1	101.2		
3		100.3	101.5		
4		98.3	99.7		
5		100.4	101.6		
6		99.4	100.9		
1		98.7	99.3		
2		98.6	99.3		
3	Intermediate precision	98.4	99.2		
4	memediate precision	98.6	99.4		
5		98.6	99.2		
6		98.6	99.2		
	Mean	99.1	100.1		
	SD	0.76	0.98		
	%RSD	0.77	0.98		



ANALYTICAL METHOD VALIDATION REPORT FOR ASSAY AND UNIFORMITY OF CONTENT OF RAMITHIAZIDE (RAMIPRIL AND HYDROCHLOROTHIAZIDE TABLETS) BY HPLC METHOD

Report Number QBD-REP-332 **Page No.** 35 of 49

Conclusion:

- 1. The relative standard deviation for peak area of Ramipril and Hydrochlorothiazide of standard solution is within acceptance criteria.
- 2. The tailing factor for Ramipril and Hydrochlorothiazide peak in standard solution is within acceptance criteria.
- 3. The theoretical plates for Ramipril and Hydrochlorothiazide peak in standard solution is within acceptance criteria.
- 4. % Assay value of Ramipril and Hydrochlorothiazide for sample solution is within specified limit.
- 5. % RSD of Assay value of Ramipril and Hydrochlorothiazide for six sample solution is within acceptance criteria.
- 6. The cumulative RSD for % Assay of Ramipril and Hydrochlorothiazide for twelve samples in method precision and intermediate precision is within acceptance criteria.

II) Intermediate precision for Content Uniformity:

Acceptance criteria:

- 1. The relative standard deviation for peak area of Ramipril and Hydrochlorothiazide of standard solution should be not more than 2.0%
- 2. The tailing factor for Ramipril and Hydrochlorothiazide peak in standard solution should be not more than 2.0
- 3. The theoretical plates for Ramipril and Hydrochlorothiazide peak in standard solution should be not less than 2000
- 4. The acceptance value of Ramipril and Hydrochlorothiazide for ten sample solution should be not more than 15.0
- % Difference of mean of content uniformity of Ramipril and Hydrochlorothiazide for Method precision and Intermediate precision should be not more than 3.0%



ANALYTICAL METHOD VALIDATION REPORT FOR ASSAY AND UNIFORMITY OF CONTENT OF RAMITHIAZIDE (RAMIPRIL AND HYDROCHLOROTHIAZIDE TABLETS) BY HPLC METHOD

Report Number	QBD-REP-332	Page No.	36 of 49

Standard details (Ramipril 5 mg & Hydrochlorothiazide 12.5 mg & Ramipril 10 mg & Hydrochlorothiazide 25 mg)

Injection No.	Retention Time	Area of Hydrochlorothia Standard	zide	Retei Tii		Area of Ramipril Standard
1	2.107	26757.552		3.9	73	7000.661
2	2.110	26742.164		3.9	77	7017.555
3	2.107	26729.583		3.9	74	7006.084
4	2.112	26720.862		3.9	72	7009.540
5	2.107	26715.221		3.9	73	7010.196
6	2.112	26708.079		3.972		6990.409
Average	2.109	26728.910		3.9	74	7005.741
SD	0.00	18.34		0.0	0	9.33
%RSD	0.12	0.07		0.05		0.13
,		System Suitabi	lity R	esults		
% RSD of Area	a of Hydrochl	orothiazide	_	.07	NMT 2.0%	
% RSD of Area of Ramipril			0.	.13	NMT 2.0%	
Tailing factor of Hydrochlorothiazide			1.	.08	NMT 2.0	
Tailing factor of Ramipril			1.	03	NMT 2.0	
Theoretical plates of Hydrochlorothiazide			10	583		NLT 2000
Theoretical plates of Ramipril			11	081	NLT 2000	

Standard details (Ramipril 10 mg & Hydrochlorothiazide 12.5 mg)

Injection No.	Retention Time	Area of Hydrochlorothiazide Standard		Reten Tin		Area of Ramipril Standard	
1	2.118	13481.003		3.944		7266.084	
2	2.115	13300.734		3.94	12	7175.544	
3	2.113	13314.339		3.94	16	7203.315	
4	2.118	13327.831		3.94	14	7248.287	
5	2.117	13325.621		3.94	14	7168.018	
6	2.118	13327.657		3.945		7219.029	
Average	2.117	13346.198		3.94	14	7213.380	
SD	0.00	66.87		0.00		39.04	
%RSD	0.10	0.50		0.03		0.54	
·		System Suitabi	lity R	esults			
% RSD of Area	of Hydrochl	orothiazide	0.	50	NMT 2.0%		
% RSD of Area of Ramipril			0.	54	NMT 2.0%		
Tailing factor of Hydrochlorothiazide			1.	13	NMT 2.0		
Tailing factor of Ramipril				04	NMT 2.0		
Theoretical plates of Hydrochlorothiazide				65	NLT 2000		
Theoretical plates of Ramipril				71	NLT 2000		



ANALYTICAL METHOD VALIDATION REPORT FOR ASSAY AND UNIFORMITY OF CONTENT OF RAMITHIAZIDE (RAMIPRIL AND HYDROCHLOROTHIAZIDE TABLETS) BY HPLC METHOD

Report Number	QBD-REP-332	Page No.	37 of 49
---------------	-------------	----------	----------

i) Intermediate precision sample details for Hydrochlorothiazide (Ramipril 5 mg & Hydrochlorothiazide 12.5 mg):

Sample	Tablets Weight (mg)	Peak Area of Sample	% Content
1	199.98	26738.909	101.7
2	195.70	26777.155	101.8
3	199.17	26455.656	100.6
4	194.89	26811.343	101.9
5	196.42	26641.010	101.3
6	197.84	26754.995	101.7
7	195.91	26962.827	102.5
8	197.34	26777.232	101.8
9	197.05	26783.011	101.8
10	197.94	26719.267	101.6
		Min	100.6
		Max	102.5
		Mean	101.7
		SD	0.49
		% RSD	0.48
		AV Value	1.4

ii) Intermediate precision sample details for Ramipril (Ramipril 5 mg & Hydrochlorothiazide 12.5 mg):

Sample	Tablets Weight (mg)	Peak Area of Sample	% Content
1	199.98	6811.163	100.7
2	195.70	6536.646	96.7
3	199.17	6440.453	95.2
4	194.89	6638.071	98.2
5	196.42	6489.863	96.0
6	197.84	6595.169	97.5
7	195.91	6659.570	98.5
8	197.34	6602.860	97.6
9	197.05	6587.300	97.4
10	197.94	6578.018	97.3
		Min	95.2
		Max	100.7
		Mean	97.5
		SD	1.49
		% RSD	1.53
		AV Value	4.6



ANALYTICAL METHOD VALIDATION REPORT FOR ASSAY AND UNIFORMITY OF CONTENT OF RAMITHIAZIDE (RAMIPRIL AND HYDROCHLOROTHIAZIDE TABLETS) BY HPLC METHOD

Report Number	QBD-REP-332	Page No.	38 of 49

iii)Intermediate precision sample details for Hydrochlorothiazide (Ramipril 10 mg & Hydrochlorothiazide 12.5 mg):

Sample	Tablets Weight (mg)	Peak Area of Sample	% Content
1	195.98	13502.588	99.0
2	201.22	14463.008	106.0
3	195.81	13649.699	100.0
4	200.20	13756.050	100.8
5	195.30	13677.799	100.2
6	199.39	13619.010	99.8
7	197.59	13654.006	100.1
8	199.74	13587.821	99.6
9	198.10	13745.397	100.7
10	200.49	13510.451	99.0
		Min	99.0
		Max	106.0
		Mean	100.5
		SD	2.02
		% RSD	2.01
		AV Value	6.0

iv) Intermediate precision sample details for Ramipril (Ramipril 10 mg & Hydrochlorothiazide 12.5 mg):

Sample	Tablets Weight (mg)	Peak Area of Sample	% Content
1	195.98	7096.170	101.7
2	201.22	7127.199	102.2
3	195.81	7127.021	102.2
4	200.20	7085.981	101.6
5	195.30	7018.480	100.6
6	199.39	6980.559	100.1
7	197.59	7081.134	101.5
8	199.74	7101.660	101.8
9	198.10	6986.968	100.1
10	200.49	7089.246	101.6
		Min	100.1
		Max	102.2
		Mean	101.3
		SD	0.78
		% RSD	0.77
		AV Value	1.9



ANALYTICAL METHOD VALIDATION REPORT FOR ASSAY AND UNIFORMITY OF CONTENT OF RAMITHIAZIDE (RAMIPRIL AND HYDROCHLOROTHIAZIDE TABLETS) BY HPLC METHOD

	Report Number	QBD-REP-332	Page No.	39 of 49	_
_			1 0.50 1 10.	0,01,	

v) Intermediate precision sample details for Hydrochlorothiazide (Ramipril 10 mg & Hydrochlorothiazide 25 mg):

Sample	Tablets Weight (mg)	Peak Area of Sample	% Content
1	199.97	25578.397	95.5
2	201.56	26562.750	99.2
3	197.78	26813.157	100.1
4	195.75	27476.889	102.6
5	200.69	26955.541	100.7
6	199.87	28934.875	108.1
7	199.91	26756.172	99.9
8	198.52	26881.100	100.4
9	196.46	27172.322	101.5
10	202.30	26817.056	100.1
		Min	95.5
		Max	108.1
		Mean	100.8
		SD	3.14
		% RSD	3.11
		AV Value	7.5

vi) Intermediate precision sample details for Ramipril (Ramipril 10 mg & Hydrochlorothiazide 25.5 mg):

Sample	Tablets Weight (mg)	Peak Area of Sample	% Content
1	199.97	6379.967	91.9
2	201.56	6778.099	97.6
3	197.78	6965.385	100.3
4	195.75	6849.165	98.6
5	200.69	6779.973	97.6
6	199.87	7308.197	105.2
7	199.91	6841.221	98.5
8	198.52	6855.122	98.7
9	196.46	6839.633	98.5
10	202.30	6756.875	97.3
		Min	91.88
		Max	105.24
		Mean	98.4
		SD	3.26
		% RSD	3.31
		AV Value	7.9



ANALYTICAL METHOD VALIDATION REPORT FOR ASSAY AND UNIFORMITY OF CONTENT OF RAMITHIAZIDE (RAMIPRIL AND HYDROCHLOROTHIAZIDE TABLETS) BY HPLC METHOD

Report Number	QBD-REP-332	Page No.	40 of 49	
---------------	-------------	----------	----------	--

Comparision of content uniformity:

i) For Hydrochlorothiazide:

Sr.No.	Lable claim (mg)	Test	% Mean Content	% mean Difference of content	
1	Ramithiazide tablets 5mg/12.5mg	Precision	103.8	2.1	
	Kamunazide tablets 5mg/12.5mg	Intermediate precision	101.7	2.1	
2	Ramithiazide tablets 10mg/12.5mg	Precision	103.4	2.0	
	realitimazide tablets follig/12.5mg	Intermediate precision	100.5	2.9	
3	Ramithiazide tablets 10mg/25mg	Precision	102.2	1.4	
3	Kamunazide tablets 10mg/25mg	Intermediate precision	100.8	1.4	

ii) For Ramipril:

Sr. No.	Lable claim (mg)	Test	% Mean Content	% mean Difference of content	
1	Ramithiazide tablets 5mg/12.5mg	Precision	98.3	0.0	
	Tamanaziae tablets 5mg/12.5mg	Intermediate precision	97.5	0.8	
2	Ramithiazide tablets 10mg/12.5mg	Precision	101.0	0.4	
	Ramanazide tablets Tollig/12.5mg	Intermediate precision	101.3	0.4	
3	Ramithiazide tablets 10mg/25mg	Precision	97.4	1 1	
3	Kammazide tablets Tomg/25mg	Intermediate precision	98.4	1.1	

Conclusion:

- 1. The relative standard deviation for peak area of Ramipril and Hydrochlorothiazide of standard solution is within acceptance criteria.
- 2. The tailing factor for Ramipril and Hydrochlorothiazide peak in standard solution is within acceptance criteria.
- 3. The theoretical plates for Ramipril and Hydrochlorothiazide peak in standard solution is within acceptance criteria.
- 4. The acceptance value of Ramipril and Hydrochlorothiazide for ten sample solution is within acceptance criteria.
- 5. % Difference of mean of content uniformity of Ramipril and Hydrochlorothiazide for Method precision and Intermediate precision is within acceptance criteria.



ANALYTICAL METHOD VALIDATION REPORT FOR ASSAY AND UNIFORMITY OF CONTENT OF RAMITHIAZIDE (RAMIPRIL AND HYDROCHLOROTHIAZIDE TABLETS) BY HPLC METHOD

Report Number QBD-REP-332 Page No. 41 of 49

10.5. ACCURACY (RECOVERY)

The accuracy of an analytical procedure expresses the closeness of agreement between the value, which is accepted either as conventional true value or an accepted reference value and found value.

Acceptance Criteria:

- 1. The relative standard deviation for peak area of Ramipril and Hydrochlorothiazide of standard solution should be not more than 2.0%
- 2. The tailing factor for Ramipril and Hydrochlorothiazide peak in standard solution should be not more than 2.0
- 3. The theoretical plates for Ramipril and Hydrochlorothiazide peak in standard solution should be not less than 2000
- 4. % Recovery at each accuracy level should be between 98.0% and 102.0%.
- 5. Mean % Recovery of all level should be between 98.0% and 102.0%.

Accuracy for Ramipril 5 mg & Hydrochlorothiazide 12.5 mg and Ramipril 10 mg & Hydrochlorothiazide 25 mg

Accuracy of Hydrochlorothiazide:

Accuracy level (%)	Sample	Placebo weight (mg)	Amount added(ppm)	Area	Amount recovered (ppm)	% Recovery	Mean % Recovery	% RSD
	Sample 1	468.98	26.29	7354.546	26.65	101.4		
50	Sample 2	464.81	25.36	7050.771	25.55	100.8	101.2	0.41
	Sample 3	464.26	26.64	7464.556	27.05	101.5		0
	Sample 1	465.19	50.07	14024.138	50.82	101.5		
100	Sample 2	465.46	50.56	14206.420	51.48	101.8	101.6	0.17
	Sample 3	463.59	50.12	14046.120	50.90	101.6		0.17
	Sample 1	463.84	75.06	20957.790	75.95	101.2		
150	Sample 2	465.65	74.95	20998.029	76.10	101.5	101.3	0.18
	Sample 3	463.80	75.06	20970.367	76.00	101.3	101.5	0.10



ANALYTICAL METHOD VALIDATION REPORT FOR ASSAY AND UNIFORMITY OF CONTENT OF RAMITHIAZIDE (RAMIPRIL AND HYDROCHLOROTHIAZIDE TABLETS) BY HPLC METHOD

Report Number QBD-REP-332 **Page No.** 42 of 49

Accuracy of Ramipril:

Accuracy level (%)	Sample	Placebo weight (mg)	Amount added(ppm)	Area	Amount recovered (ppm)	% Recovery	Mean % Recovery	% RSD
	Sample 1	468.98	10.67	4897.906	10.79	101.1		
50	Sample 2	464.81	10.50	4799.096	10.57	100.6	101.1	0.51
	Sample 3	464.26	10.71	4944.512	10.89	101.7		
	Sample 1	465.19	19.82	8820.949	19.43	98.1		
100	Sample 2	465.46	19.86	8925.371	19.66	99.0	98.5	0.50
	Sample 3	463.59	19.83	8855.191	19.51	98.4		
	Sample 1	463.84	30.22	13783.487	30.36	100.5		
150	Sample 2	465.65	30.08	13782.042	30.36	100.9	100.7	0.22
	Sample 3	463.80	30.14	13787.558	30.37	100.7		

Accuracy for Ramipril 10 mg & Hydrochlorothiazide 12.5 mg:

Accuracy of Hydrochlorothiazide:

Accuracy level (%)	Sample	Placebo weight (mg)	Amount added(ppm)	Area	Amount recovered (ppm)	% Recovery	Mean % Recovery	% RSD
	Sample 1	467.98	12.69	3507.590	12.57	99.0		
50	Sample 2	465.83	12.70	3523.651	12.62	99.4	99.4	0.31
	Sample 3	466.32	12.66	3522.441	12.62	99.7		
	Sample 1	464.60	25.29	7032.703	25.20	99.6		
100	Sample 2	465.01	25.19	7014.152	25.13	99.8	99.7	0.07
	Sample 3	465.13	25.24	7029.037	25.18	99.8		
	Sample 1	464.49	37.56	10364.95	37.13	98.9		
150	Sample 2	466.10	37.50	10418.45	37.33	99.5	99.3	0.36
	Sample 3	464.27	37.46	10393.90	37.24	99.4		



ANALYTICAL METHOD VALIDATION REPORT FOR ASSAY AND UNIFORMITY OF CONTENT OF RAMITHIAZIDE (RAMIPRIL AND HYDROCHLOROTHIAZIDE TABLETS) BY HPLC METHOD

Report Number	QBD-REP-332	Page No.	43 of 49
---------------	-------------	----------	----------

Accuracy of Ramipril:

Accuracy level (%)	Sample	Placebo weight (mg)	Amount added(ppm)	Area	Amount recovered (ppm)	% Recovery	Mean % Recovery	% RSD
	Sample 1	467.98	10.30	4774.975	10.47	101.6		
50	Sample 2	465.83	10.19	4701.765	10.31	101.1	101.5	0.31
	Sample 3	466.32	10.36	4807.576	10.54	101.7		
	Sample 1	464.60	20.06	9199.755	20.17	100.6		
100	Sample 2	465.01	20.11	9192.101	20.15	100.2	100.4	0.18
	Sample 3	465.13	20.08	9191.341	20.15	100.3		
	Sample 1	464.49	30.06	13497.348	29.59	98.4		
150	Sample 2	466.10	30.05	13552.427	29.71	98.9	98.5	0.36
	Sample 3	464.27	30.10	13483.137	29.56	98.2		

Conclusion:

- 1. The relative standard deviation for peak area of Ramipril and Hydrochlorothiazide of standard solution is within acceptance criteria.
- 2. The tailing factor for Ramipril and Hydrochlorothiazide peak in standard solution is within acceptance criteria.
- 3. The theoretical plates for Ramipril and Hydrochlorothiazide peak in standard solution is within acceptance criteria.
- 4. % Recovery at each accuracy level is within acceptance criteria.
- 5. Mean % Recovery of all level is within acceptance criteria.

10.6. ROBUSTNESS

The robustness of an analytical procedure is a measure of its capacity to remain unaffected by small, but deliberate variations in method parameters and provides an indication of its reliability during normal usage.

Acceptance criteria:

1. The relative standard deviation for peak area of Ramipril and Hydrochlorothiazide of standard solution should be not more than 2.0%



ANALYTICAL METHOD VALIDATION REPORT FOR ASSAY AND UNIFORMITY OF CONTENT OF RAMITHIAZIDE (RAMIPRIL AND HYDROCHLOROTHIAZIDE TABLETS) BY HPLC METHOD

Report Number QBD-REP-332 Page No. | 44 of 49

- 2. The tailing factor for Ramipril and Hydrochlorothiazide peak in standard solution should be not more than 2.0
- 3. The theoretical plates for Ramipril and Hydrochlorothiazide peak in standard solution should be not less than 2000
- 4. % Difference for % Assay of sample solution in robustness and % Assay of sample solution in method precision should be not more than 2.0%

System suitability details:

Robustness parameter	% RSD of Ramipril	% RSD of HCTZ	Therotical plates of Ramipril	Therotical plates of HCTZ	Tailing factor of Ramipril	Tailing factor of HCTZ			
, , , , , , , , , , , , , , , , , , ,	Limit								
	NMT 2.0%	NMT 2.0%	NLT 2000	NLT 2000	NMT 2.0	NMT 2.0			
Precision	0.13	0.07	10410	10705	1.00	1.04			
Flow Rate 0.9 mL/min	0.16	0.11	11245	11488	0.96	1.02			
Flow Rate 1.1 mL/min	0.22	0.15	10098	10341	0.95	1.03			
Organic content decrease	0.40	0.11	5847	3881	1.13	1.30			
Organic content increase	0.66	0.18	5015	3757	1.16	1.26			
Buffer pH 2.3	0.21	0.22	4532	3390	1.07	1.19			
Buffer pH 2.7	0.24	0.11	5019	3475	1.10	1.19			

Robustness study details:

For Hydrochlorothiazide:

Robustness parameter	% Assay	% Difference (NMT 2.0 %)
Precision	99.6	NA
Flow Rate 0.9 mL/min	100.1	0.5
Flow Rate 1.1 mL/min	99.9	0.3
Organic content decrease	98.9	0.7
Organic content increase	98.4	1.2
Buffer pH 2.3	98.4	1.2
Buffer pH 2.7	98.3	1.3



ANALYTICAL METHOD VALIDATION REPORT FOR ASSAY AND UNIFORMITY OF CONTENT OF RAMITHIAZIDE (RAMIPRIL AND HYDROCHLOROTHIAZIDE TABLETS) BY HPLC METHOD

Report Number	QBD-REP-332	Page No.	45 of 49
---------------	-------------	----------	----------

Robustness study details:

For Ramipril:

Robustness parameter	% Assay	% Difference (NMT 2.0 %)
Precision	100.9	NA
Flow Rate 0.9 mL/min	100.9	0.0
Flow Rate 1.1 mL/min	100.8	0.1
Organic content decrease	100.1	0.8
Organic content increase	100.7	0.2
Buffer pH 2.3	100.0	0.9
Buffer pH 2.7	101.7	0.8

Conclusion:

- 1. The relative standard deviation for peak area of Ramipril and Hydrochlorothiazide of standard solution is within acceptance criteria.
- 2. The tailing factor for Ramipril and Hydrochlorothiazide peak in standard solution is within accepta-ce criteria.
- 3. The theoretical plates for Ramipril and Hydrochlorothiazide peak in standard solution is within acceptance criteria.
- 4. % Difference for % Assay of sample solution in robustness and % Assay of sample solution in method precision is within acceptance criteria.

10.7. FILTER STUDY

Acceptance criteria:

- 1. The relative standard deviation for peak area of Ramipril and Hydrochlorothiazide of standard solution should be not more than 2.0%
- 2. The tailing factor for Ramipril and Hydrochlorothiazide peak in standard solution should be not more than 2.0
- 3. The theoretical plates for Ramipril and Hydrochlorothiazide peak in standard solution should be not less than 2000
- 4. %Difference between filtered solution through different filters and centrifuged sample solution should be NMT 2.0%



ANALYTICAL METHOD VALIDATION REPORT FOR ASSAY AND UNIFORMITY OF CONTENT OF RAMITHIAZIDE (RAMIPRIL AND HYDROCHLOROTHIAZIDE TABLETS) BY HPLC METHOD

Report Number	QBD-REP-332	Page No.	46 of 49
To the state of th	QDD 14E1 332	Tage 110.	40 01 49

Standard details (Ramipril 5 mg & Hydrochlorothiazide 12.5 mg & Ramipril 5 mg & Hydrochlorothiazide 12.5 mg)

Injection No.	Retention Time	Area of Hydrochlorothia Standard	zide	Reter Tir		Area of Ramipril Standard
1	2.225	29779.744		3.8	82	6982.083
2	2.225	29910.378		3.8	83	6990.418
3	2.227	29895.007		3.8	85	6982.654
4	2.227	29870.134		3.8	85	6978.374
5	2.225	29937.184		3.8	83	6982.466
6	2.225	29970.544		3.883		6990.327
Average	2.226	29893.832		3.8	84	6984.387
SD	0.001	65.75		0.0	01	4.89
%RSD	0.05	0.22	0.0		3	0.07
		System Suitabi	lity R	esults		
% RSD of Area		orothiazide	0.	.22		NMT 2.0%
% RSD of Area			0.	.07		NMT 2.0%
Tailing factor of Hydrochlorothiazide		1.	15		NMT 2.0	
Tailing factor of Ramipril		1.	06	NMT 2.0		
Theoretical plates of Hydrochlorothiazide		60	940			
Theoretical plan	tes of Ramipr	il	85	84		NLT 2000

Filter study details:

For Hydrochlorothiazide:

Robustness parameter	% Assay	% Difference (NMT 2.0 %)
Centrifuged	100.8	-
Filtered through 0.22 u filter	100.8	0.02
Filtered through 0.45 u filter	100.9	0.15

Filter study details:

For Ramipril:

Robustness parameter	% Assay	% Difference (NMT 2.0 %)
Centrifuged	101.5	-
Filtered through 0.22 u filter	101.5	0.08
Filtered through 0.22 u filter	101.5	0.03



ANALYTICAL METHOD VALIDATION REPORT FOR ASSAY AND UNIFORMITY OF CONTENT OF RAMITHIAZIDE (RAMIPRIL AND HYDROCHLOROTHIAZIDE TABLETS) BY HPLC METHOD

Report Number QBD-REP-332	Page No.	47 of 49
---------------------------	----------	----------

Conclusion:

- 1. The relative standard deviation for peak area of Ramipril and Hydrochlorothiazide of standard solution is within acceptance criteria.
- 2. The tailing factor for Ramipril and Hydrochlorothiazide peak in standard solution is within acceptance criteria.
- 3. The theoretical plates for Ramipril and Hydrochlorothiazide peak in standard solution is within acceptance criteria.
- 4. %Difference between filtered solution through different filters and centrifuged sample solution is within acceptance criteria.

10.8. SOLUTION STABILITY

Acceptance criteria:

Standard solution stability:

The RSD of area response of repeatedly injected standard solution over a period of 24 hrs should be not more than 2.0%

Sample solution stability:

The RSD of area response of repeatedly injected sample solution over a period of 24 hrs should be not more than 2.0%

Standard details (Ramipril 5 mg & Hydrochlorothiazide 12.5 mg & Ramipril 5 mg & Hydrochlorothiazide 12.5 mg)

Injection Retention Time 1 2.097		Area of Hydrochlorothiazide Standard	Retention Time	Area of Ramipril Standard
		27140.043	3.650	6928.843
2 2.097		27182.012	3.650	6943.845
3 2.097		27163.638	3.650	6935.208
4	2.097	27160.989	3.650	6940.622
5	2.097	27129.962	3.650	6923.226
6 2.097		27176.337	3.650	6945.313
Average 2.097		27158.830	3.650	6936.176
SD	0.000	20.29	0.000	8.77
%RSD	0.00	0.07	0.00	0.13



ANALYTICAL METHOD VALIDATION REPORT FOR ASSAY AND UNIFORMITY OF CONTENT OF RAMITHIAZIDE (RAMIPRIL AND HYDROCHLOROTHIAZIDE TABLETS) BY HPLC METHOD

Report Number QBD-REP-332	Page No.	48 of 49	
----------------------------------	----------	----------	--

System Suitability Results							
% RSD of Area of Hydrochlorothiazide	0.07	NMT 2.0%					
% RSD of Area of Ramipril	0.00	NMT 2.0%					
Tailing factor of Hydrochlorothiazide	1.04	NMT 2.0					
Tailing factor of Ramipril	1.00	NMT 2.0					
Theoretical plates of Hydrochlorothiazide	10705	NLT 2000					
Theoretical plates of Ramipril	10410	NLT 2000					

Solution stability standard:

Time (hr)	Area of Standard for HCTZ	Area of Standard for Ramipril	%RSD (NMT 2.0%) HCTZ	%RSD (NMT 2.0%) Ramipril
Initial	27140.044	6928.843	NA	NA
3 hrs	27272.086	6969.910	0.34	0.42
5 hrs	27477.861	7028.509	0.62	0.72
13 hrs	27578.480	7048.360	0.72	0.78
19 hrs	27900.150	7159.078	1.07	1.25
25 hrs	28227.280	7213.154	1.47	1.55
28 hrs	28438.752	7265.373	1.76	1.79

Sample solution stability details:

Sample details	Area of Sample		% 1	Assay	%RSD (NMT 2.0%)	%RSD (NMT 2.0%)
	HCTZ	Ramipril	HCTZ	Ramipril	HCTZ	Ramipril
Initial	26811.141	6690.018	98.8	98.6	NA	NA
4 hrs	26774.333 6684.338 26925.700 6714.042		98.7	98.7 98.5		0.08
7hrs			99.2	99.0	0.42	0.35
10 hrs	26949.482	6741.319	99.3	99.4	0.51	0.76
16 hrs	27177.318	6797.521	100.2	100.2	1.35	1.58
22 hrs	27285.001 6819.047		100.6	100.5	1.75	1.90
25 hrs	27424.241	6854.231	101.1	101.0	2.26	2.42

Conclusion:

Standard solution stability:

The standard solution is stable upto 28hrs.



ANALYTICAL METHOD VALIDATION REPORT FOR ASSAY AND UNIFORMITY OF CONTENT OF RAMITHIAZIDE (RAMIPRIL AND HYDROCHLOROTHIAZIDE TABLETS) BY HPLC METHOD

Report Number QBD-REP-332 Page No. 49 of 49

Sample solution stability:

Sample solution is stable upto 22 hrs.

10 DOCUMENTATION

All validation activities shall be recorded online in to the laboratory notebook. At the end of the Validation, a report shall be generated. All results will be discussed, conclusions drawn and any deviations from the protocol will be documented in the report.

11 REFERENCES

ICH guidelines

12 REVISION HISTORY

Version	Date	Changes	Reason for change	
0		Not Applicable	New	

INVOICE

QbD Research And Development Lab Pvt Ltd

2nd Floor, TSSIA House Plot No. P-26, Road No.16 / T Wagle Ind Estate, Thane 400604 CIN No.U73100MH2012PTC232153 GSTIN/UIN: 27AAACQ2815J1ZG

E-Mail: lab@qbdresearch.com

Generic Healthcare Pvt Ltd

City Squre,29/A,Office No- 205, 2nd Floor, Behind Hotel Pride, Shivaji Nager, Pune-, 411005, K/A Mr Palve Sir,

State Name GSTIN/UIN

: Maharashtra, Code : 27 : 27AADCG9221D1ZU

Invoice No.

Dated

QbD/117/2017-18

27-Jul-2017

Supplier's Ref.

Other Reference(s)

Buyer's Order No.

Dated

SI No		Quantity	Rate	per	Disc. %	Amount
1	Ramithiazide Tablet Analytical Method Validation For Assay and Uniformity of Content Of Ramithiazide by HPLC Method	1 No.	4,00,000.00	No.		4,00,000.00
		ST Output ST Output	-	%		36,000.00 36,000.00

Total 1 No.

₹ 4,72,000.00

Amount Chargeable (in words)

E. & O.E

Indian Rupees Four Lakh Seventy Two Thousand Only

	HSN/SAC		Taxable	Cent	tral Tax	Sta	te Tax	Total
			Value	Rate	Amount	Rate	Amount	Tax Amount
998346			4,00,000.00	9%	36,000.00	9%	36,000.00	72,000.00
		Total	4,00,000.00		36,000.00		36,000.00	72,000.00

Tax Amount (in words): Indian Rupees Seventy Two Thousand Only

Company's Bank Details

Bank Name

A/c No.

TJSB Sahakari Bank Ltd-CC-47

Branch & IFS Code: Mulund (W) & TJSB00000

CC A/c No.01313010000004

for QbD Research And Deve

Company's PAN

: AAACQ2815J

Declaration

1.Payment Terms : Payment should be me made within 10 days on recipt of invoice copy.

2 Interest at 21% will be charged on this bill if not paid within one month.

This is a Computer Generated Invoice