

	FINISHED PRODUCT SPECIFICATION		Page No.	1 of 2
	Quality Control Department			
Product Name	NOVOMIX GEN II-65051			
Colour	Pink			
Specification No.	FP/NOVOMPS/C-1/GEN II-65051/01	Supersedes No.	NA	
Customer's Name	MSN Laboratories			
Reference	In House	Effective Date	12/06/2024	
Ref. Annexure No.	A/SOP/QC/030/04	Review Month	May - 2029	

GENERAL INFORMATION




Pharmacopoeial reference : In House

Quantity to be sampled :

Analysis Sample	Control sample
25 g	75 g

Shelf Life : 24 Months

Storage : Store in dry place, in well closed container, away from direct heat and light.

	PREPARED BY	REVIEWED BY	APPROVED BY
Name	Rajanikant Dandekar	Manisha Dhole	Rahul Bankar
Sign/Date	 12/06/24	 12/06/24	 12/06/24
Designation	Sr. Q.C. Executive	Sr. Executive	Sr. Executive
Department	Quality Control	Quality Control	Quality Assurance

	FINISHED PRODUCT SPECIFICATION	Page No.	2 of 2
	Quality Control Department		
Product Name	NOVOMIX GEN II-65051		
Specification No.	FP/NOVOMPS/C-1/GEN II-65051/01		




Qualitative Formula:

1. Hydroxypropyl methyl cellulose USP
2. Polyethylene Glycol USP
3. Lactose Monohydrate USP
4. Titanium Dioxide USP
5. Yellow Iron Oxide USP
6. Red Iron Oxide USP

Sr. No.	Test	Specification	Reference
1.	Appearance	Pink powder	GTP/QC/001
2.	Identification by IR	IR spectrum of test sample should be concordant with the IR spectrum of standard.	GTP/QC/009
3.	Colour Difference		GTP/QC/008
	1) DE (By colour spectrophotometer)	NMT 1.5	
	2) Visually	Colour shade should match visually with Standard	
4.	Dispersion	Pass	GTP/QC/006
5.	Ash Content	22.02 % w/w to 29.79 % w/w	GTP/QC/004




REVISION HISTORY FOR SPECIFICATION:

Revision No.	Revision Description	Effective Date
01	New Document	12/06/2024

	PREPARED BY	REVIEWED BY	APPROVED BY
Sign/Date			

	FINISHED PRODUCT STANDARD TEST PROCEDURE (STP)		Page No.	1 of 4
	Quality Control Department			
Product Name	NOVOMIX GEN II-65051			
Colour	Pink			
STP No.	FP/NOVOMSTP/C-1/GEN II-65051/01			
Reference	In House	Supersedes No.	NA	
Customer's Name	MSN Laboratories			
Ref. SPEC No.	FP/NOVOMPS/C-1/GEN II-65051/01	Effective Date	12/06/2024	
Ref. Annexure No.	A/SOP/QC/030/08	Review Month	May-2023	

01.	Appearance : Instrument: Balance Take about 2 to 3 g of test sample and transferred into a clean and dry petri dish. Spread it uniformly by spatula and observe. Specification: Pink powder.
02.	Identification by IR: Instrument: Balance, IR Spectrophotometer, Oven Take about 1 mg of sample to be analyzed and 100 mg IR grade KBr onto piece of weighing paper. The KBr should be store in the desiccator when not in used. Transfer the sample /KBr mixture to the agate mortar. Thoroughly disperse the mixture using the spatula or spoon. The pestle is then used with a grinding motion to homogenously disperse the mixture component. Completed homogeneity takes approximately 1-2 minutes of grinding the components with the pestle. Prepare a KBr pellet using an appropriate apparatus (Press Kit, E- Z press etc.) carefully examine the pallet film to ensure that it is intact and somewhat transparent. Run the pallet on suitable spectrophotometer from 4000 cm^{-1} to 650 cm^{-1} according to the operating instruments. All peak should be match the reference with relation to wavelength location and approximate magnitude location. Overlap the sample spectra with reference spectra. Note: Sample and KBr to be dried at 105°C for about 30 minutes before use Or

	PREPARED BY	REVIEWED BY	APPROVED BY
Name	Rajanikant Dudhkar	Manisha Dhere	Rahul Bankar
Sign/Date	 12/06/24	 12/06/24	 12/06/24
Designation	Sr. Q.C. Executive	Sr. Executive	Sr. Executive
Department	Quality Control	Quality Control	Quality Assurance

	FINISHED PRODUCT STANDARD TEST PROCEDURE (STP)		Page No.	2 of 4
	Quality Control Department			
Product Name:	NOVOMIX GEN II-65051			
STP No. :	FP/NOVOMSTP/C-1/GEN II-65051/01			

By ATR Accessory:

Make sure the crystal is free from any residue by wiping clean with IPA or methanol and delicate wipe. Perform the background measurement. Grind a small amount of sample using agate mortar. Transfer enough of the undiluted sample to completely cover the crystal. Lower the ATR press all the way down with sufficient pressure to ensure a well-defined scan. Collect a spectrum of sample from 4000 cm⁻¹ to 650 cm⁻¹.

All peak should be match the reference with relation to wavelength location and approximate magnitude location. Overlap the sample spectra with reference spectra.

Specification: IR spectrum of test sample should be concordant with the IR spectrum of standard.

03. Colour difference :

Instrument : Balance , Applicator, Colour matching spectrophotometer

1) Visually: Transfer mixture of 20 ml of dichloromethane and 10 ml methanol into a suitable beaker. Then lower the paddle into the solvent mixture and switch on the stirrer. Adjust the speed of the stirrer so that a vortex is produced without pulling in any air. Check that the head of stirrer is turning without any 'wobble'. Then add 2.0 g of test sample as quickly as the stirrer will allow without excess build-up of product on the surface. The speed of the stirrer will need to be increased to maintain the vortex but must remain in the operating range of 800 to 1200 rpm. Mix for 20-30 minutes. After mixing pass the sample solution through 100 mesh.

Instrument Method parameters: Speed: 10 mm/sec.

Distance : 50%

Thickness: 800 µm




Volume : About 4 ml

Drying Time : 10-15 minutes

Drying Temp.: Room Temperature




Place a labeled card (Ivory or equivalent) on the drawdown plate with the matte side facing up and apply a vacuum. Prepare colour shade card using above method.

Note: Dispersion media and sample quantity may vary as per requirement but final concentration

	PREPARED BY	REVIEWED BY	APPROVED BY
Sign/Date			

	FINISHED PRODUCT STANDARD TEST PROCEDURE (STP)		Page No.	3 of 4
	Quality Control Department			
Product Name:	NOVOMIX GEN II-65051			
STP No. :	FP/NOVOMSTP/C-1/GEN II-65051/01			




	should remain same. Compare the colour shade cards with a standard reference colour shade card. Specification: Colour shade should match visually with Standard. 2) DE (By colour spectrophotometer): Compare the colour of sample shade cards with a reference standard already stored in the colour matching system. Specification: NMT 1.5
04.	Dispersion: Instrument: Balance Procedure: Arrange the sieves as 30 mesh with receiver. Weigh accurately 10 g of sample. Sieve through the mesh without applying pressure. No retained material remains on sieve. Specification: Should Pass
05.	Ash Content : Instrument: Balance, Muffle Furnace Heat a silica crucible at $600 \pm 25^{\circ} \text{C}$ for 15 minutes, allow to cool in a desiccator and weigh. Take 1-2 g of sample to the crucible and weigh the crucible. Ignite the sample at $600 \pm 25^{\circ} \text{C}$ for 45 minutes. Allow the silica crucible to come to room temperature in a desiccator before weighing. Repeat the operation until two successive weighing do not differ by more than 0.5 mg. Calculation: $\% \text{ Ash} = \frac{(W4-W1)}{(W2-W1)} \times 100$ Where, W1 = Weight of the empty crucible (g) W2 = Weight of the crucible with test sample before ignition (g) W3 = Weight of the crucible with residue after ignition (g) W4= Weight of the crucible with residue after ignition (g) (Constant weight) Specification: 22.02 % w/w to 29.79 % w/w

	PREPARED BY	REVIEWED BY	APPROVED BY
Sign/Date	 12/06/24	 12/06/24	 12/06/24

	FINISHED PRODUCT STANDARD TEST PROCEDURE (STP)	Page No.	4 of 4
	Quality Control Department		
Product Name:	NOVOMIX GEN II-65051		
STP No. :	FP/NOVOMSTP/C-1/GEN II-65051/01		

REVISION HISTORY FOR STANDARD TEST PROCEDURE:

Revision No.	Revision Description	Effective Date
01	New Document	1210612024

	PREPARED BY	REVIEWED BY	APPROVED BY
Sign/Date	 12106124	 12106124	 12106124